

FIXED POINT THEOREM FOR THREE ASYMPTOTICALLY NONEXPANSIVE MAPPINGS THROUGH IMPLICIT ITERATIVE SCHEME

Anil Rajput¹, Abha Tenguria² & Sanchita Pandey³

¹Department of mathematics, CSA Govt. PG College, Nodal Sehore

²Department of mathematics, Govt. M.L.B College, Bhopal

³ Research scholar, Govt. PG College, Sehore

ABSTRACT

In this paper, a new iterative scheme for approximating common fixed point of three asymptotically nonexpansive mapping is defined and we have proved weak and strong convergence theorems in a uniformly convex Banach space. A two-step iteration scheme for asymptotically nonexpansive mappings in uniformly convex Banach space includes Ishikawa type and Mann type iterations as special cases. The results obtained in this paper represent an extension as well as refinement of previous known results.

AMS Subject Classification: 47H05, 47H10, 49M05.

Keywords: Iteration process, asymptotically nonexpansive, condition (A'), L-Lipschitzian mapping, Opial's condition.

1. INTRODUCTION

This class of asymptotically nonexpansive mappings was to introduced by Goebel and Kirk [9] in 1972. They proved that, if K is a nonempty bounded closed convex subset of a uniformly convex Banach space E , then every asymptotically nonexpansive self-mapping T of K has a fixed point. The fixed point iteration process for asymptotically nonexpansive mapping in Banach spaces including Mann and Ishikawa iterations processes have been studied extensively by many authors; see ([2]-[24]). Throughout this paper, N denotes the set of all positive integers and E be a uniformly convex Banach space, K be a nonempty closed convex subset of E and $F(T) : T x = x$. A mapping $T: K \rightarrow K$ is said to be asymptotically nonexpansive if for a sequence $\{k_n\} \subset [1, \infty)$ with $\lim_{n \rightarrow \infty} k_n = 1$, if $\|T^n x - T^n y\| \leq k_n \|x - y\|$, for all $x, y \in K$ and for all $n \in N$.

In 1991, Schu ([6], [7]) introduced a modified Mann iteration process to approximate fixed points of asymptotically nonexpansive self-map defined on nonempty closed convex and bounded subset of Hilbert space H .

In 2001, Xu and Ori [5] introduced the following implicit iteration scheme for common fixed points of a finite family of nonexpansive mappings $\{T_i\}_{i=1}^N$ in Hilbert spaces:

$$x_n = \alpha_n x_{n-1} + (1 - \alpha_n) T_n x_n, n \geq 1,$$
 where $T_n = T_n \text{ mod } N$, and they proved weak convergence theorem.

In 2008 Zhao et al. [8] introduced the following iteration scheme for common fixed points of nonexpansive mapping T in Banach space and proved weak and strong convergence theorems:

$$x_n = \alpha_n x_{n-1} + \beta_n T x_{n-1} + \gamma_n T x_n, n \geq 1,$$
 Where $\{\alpha_n\}, \{\beta_n\}$ and $\{\gamma_n\}$ are real sequences in $[0, 1]$, and $\alpha_n + \beta_n + \gamma_n = 1$.

The Picard and Mann [20] iteration schemes for a mapping $T : K \rightarrow K$ are defined by

$$\begin{cases} x_1 = x \in K, \\ x_{n+1} = T^n x_n \end{cases} \quad (1)$$

and

$$\begin{cases} x_1 = x \in K, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n, \end{cases} \quad n \in N \quad (2)$$

Where $\{\alpha_n\}$ is in $(0,1)$. It is well-known that Picard iteration scheme converges for contractions but not converges for nonexpansive mapping whereas Mann iteration scheme converges for n -onexpansive. Several authors have been studied weak and strong convergence problems of iterative sequence (with errors) for asymptotically nonexpansive type mappings in a Hilbert space or a Banachspace .(see[25],[11], [22],[6]).

In 2007, Agrawal et al. [17] introduced the following iteration process:

$$\begin{cases} x_1 = x \in K, \\ x_{n+1} = (1 - \alpha_n)T^n x_n + \alpha_n T^n y_n, \\ y_n = (1 - \beta_n)x_n + \beta_n T^n x_n, \end{cases} \quad n \in N \quad (3)$$

Where $\{\alpha_n\}$ and $\{\beta_n\}$ is in $(0,1)$. They showed that this process converges at a rate same as that of Picard iteration and faster than Mann iteration for contractions. The above process deals with one mapping only. The case of two mappings in iterative processes has also remained under study since Das and Debata [2] gave and studied a two mappings process. Also see, for example [19] and [23]. The problem of approximating common fixed points of finitely many mapping plays an important role in applied mathematics especially in the theory of evaluation equation and the minimization problems. See ([12], [13],[14], [24]) for example.

In 2001, Khan and Takahashi [19] approximated the fixed points of two asymptotically nonexpansive mappings $S, T : K \rightarrow K$ through the sequence $\{x_n\}$ given by

$$\begin{cases} x_1 = x \in K, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n S^n y_n, \\ y_n = (1 - \beta_n)x_n + \beta_n T^n x_n, \end{cases} \quad n \in N \quad (4)$$

Where $\{\alpha_n\}$ and $\{\beta_n\}$ is in $(0,1)$.

Khan et al. [18] modified the iteration process (4) to the case of two mappings as follows:

$$\begin{cases} x_1 = x \in K, \\ x_{n+1} = (1 - \alpha_n)T^n x_n + \alpha_n S^n y_n, \\ y_n = (1 - \beta_n)x_n + \beta_n T^n x_n, \end{cases} \quad n \in N \quad (5)$$

Where $\{\alpha_n\}$ and $\{\beta_n\}$ is in $(0,1)$.

Recently, Yadav et al.[15] . introduced a new implicit iteration scheme as below:

$$\begin{cases} x_1 = x \in K, \\ x_{n+1} = \alpha_n x_n + \beta_n T^n x_n + \gamma_n S^n y_n \\ y_n = \alpha'_n x_n + \beta'_n S^n x_n + \gamma'_n T^n x_n, \end{cases} \quad n \in N \quad (6)$$

Where $\{\alpha_n\}$, $\{\beta_n\}$, $\{\gamma_n\}$, $\{\alpha'_n\}$, $\{\beta'_n\}$, $\{\gamma'_n\}$, is in $[0,1]$ and $\alpha_n + \beta_n + \gamma_n = \alpha'_n + \beta'_n + \gamma'_n = 1$ for fixed points of asymptotically nonexpansive mapping T in uniformly convex Banach space.

In this if set $T=I$, $S=T$ and $\gamma_n = \alpha_n$ and $\beta'_n = 0$ then the scheme will reduce to the Mann type iteration scheme given by:

$$\begin{cases} x_1 = x \in K, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n, \end{cases} \quad n \in N \quad (7)$$

Above iteration scheme is introduced in 1991, by Schu [7] for modified Mann iteration process to approximate fixed points of asymptotically nonexpansive self-map, where $T : k \rightarrow k$ is an asymptotically nonexpansive mapping with a sequence $\{k_n\}$ such that $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ and $\{\alpha_n\}$ is a sequence in $(0,1)$ satisfying the condition $\delta \leq \alpha_n \leq 1 - \delta$ all $n \in \mathbb{N}$ and for some $\delta > 0$. Then the sequence $\{x_n\}$ converges weakly to a fixed point of T .

On this basis we have introduced new implicit iterative scheme for three mappings as follows:

$$\begin{cases} x_1 = x \in K, \\ x_{n+1} = a_n x_n + b_n T^n x_n + c_n R^n x_n + d_n S^n y_n \\ y_n = a_n' x_n + b_n' S^n x_n + c_n' T^n x_n + d_n' R^n x_n, n \in \mathbb{N} \end{cases} \tag{8}$$

Where $\{a_n\}, \{b_n\}, \{c_n\}, \{d_n\}, \{a_n'\}, \{b_n'\}, \{c_n'\}, \{d_n'\}$ is in $[0,1]$ and $a_n + b_n + c_n + d_n = a_n' + b_n' + c_n' + d_n' = 1$ for fixed points of asymptotically nonexpansive mapping T in uniformly convex Banach space.

Again if we set $R, T = I$ and $b_n' = 0$ then the scheme will reduce to the Mann type iteration scheme given by:

$$\begin{cases} x_1 = x \in K, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T^n x_n, n \in \mathbb{N} \end{cases} \tag{9}$$

2. PRELIMINARIES

Let us now gather some pre-requisites. Let $X = \{x \in E : \|x\| = 1\}$ and E^* be the dual of E . The space E has:

- (i) Gateaux differentiable norm if

$$\lim_{t \rightarrow 0} \frac{\|x+ty\| - \|x\|}{t}$$

exists for each $x, y \in E$;

- (ii) Fréchet differentiable norm (see e.g. [21]) for each x in E , the above limit exists and is attained uniformly for y in E and in this case, it is also well-known that

$$\langle h, J(x) \rangle + \frac{1}{2} \|x\|^2 \leq \frac{1}{2} \|x + h\|^2 \leq \langle h, J(x) \rangle + \frac{1}{2} \|x\|^2 + b(\|h\|)$$

for all $x, h \in E$, where J is the Fréchet derivative of the function $\frac{1}{2} \|\cdot\|^2$. $\langle \cdot, \cdot \rangle$ is the dual pairing between E and E^* , and b is an increasing function defined on

$$[0, \infty) \text{ such that } \lim_{t \rightarrow 0} \frac{b(t)}{t} = 0;$$

- (iii) Opial's condition [26] if for any sequence $\{x_n\}$ in $E, x_n \rightarrow x$ implies that

$$\limsup_{n \rightarrow \infty} \|x_n - x\| < \limsup_{n \rightarrow \infty} \|x_n - y\|$$

for all $y \in E$ with $y \neq x$.

Following are the definitions and lemma used to prove the results in the next section.

Definition 1. Let K be a nonempty closed convex subset of a uniformly convex Banach space E . A mapping $T : k \rightarrow k$ is said to be asymptotically nonexpansive on k if there exists sequence $k_n, k_n \geq 1$ with $\lim k_n = 1$, such that

$$\|T^n x - T^n y\| \leq k_n,$$

for each $x, y \in K$ and each $n \geq 1$. if $k_n = 1$, then T is known as a nonexpansive mapping. \rightarrow

Definition 2 Let E be a uniformly convex Banach space, k be a nonempty closed convex subset of E , and $T : k \rightarrow k$ be an asymptotically nonexpansive mapping. Then $I - T$ is said to be demi-closed at 0, if $x_n \rightarrow x$ converges weakly and $x_n - Tx_n \rightarrow 0$ converges strongly, then it implies that $x \in K$ and $Tx = x$.

Definition 3 [3] Suppose three mappings $S, T, R : k \rightarrow k$, where k is a subset of a normed space E , said to be satisfy Condition (A) if there exists a nondecreasing function $f : [0, \infty) \rightarrow [0, \infty)$ with $f(0)=0, f(r)>0$ for all $r \in (0, \infty)$ such that either $\|x - Sx\| \geq f(d(x, F))$ or $\|x - Tx\| \geq f(d(x, F))$ or either $\|x - Rx\| \geq f(d(x, F))$ for all $x \in K$ where $d(x, F) = \inf\{\|x - p\| : p \in F = F(S) \cap F(T) \cap F(R)\}$.

Lemma 1 (see [10], Lemma 1) .Let $\{a_n\}$, $\{b_n\}$ and $\{\delta_n\}$ be sequence of non –negative real numbers satisfying the inequilty

$a_{n+1} \leq (1+\delta_n) a_n + b_n$.
If $\sum_{n=1}^{\infty} \delta_n < \infty$ and If $\sum_{n=1}^{\infty} b_n < \infty$ then $\lim_{n \rightarrow \infty} a_n$ exists. In particular, if $\{a_n\}$ has a subsequence converging to 0, then $\lim_{n \rightarrow \infty} a_n = 0$.

Lemma 2 (see [6]) Suppose that E be a uniformly convex Banach space and $0 < p \leq t_n \leq q < 1$ for all $n \in N$. Let $\{x_n\}$, $\{a_n\}, \{y_n\}$ be two sequences of E such that $\limsup_{n \rightarrow \infty} \|x_n\| \leq r$, $\limsup_{n \rightarrow \infty} \|y_n\| \leq r$ and $\lim_{n \rightarrow \infty} \|t_n x_n + (1 - t_n)y_n\| = r$ hold for some $r \geq 0$. Then $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$.

Lemma 3 Let K be a normed linear space and K be a nonempty convex subset of X . Let $T : k \rightarrow k$ be an asymptotically nonexpansive mapping with nonempty fixed point set $F(T)$ and a sequence $k_n \geq 1$ of positive real numbers such that $\sum_{n=1}^{\infty} (k_n - 1) < \infty, k_n \rightarrow 1$ as $n \rightarrow \infty$.
Let a sequence $\{x_n\}$ defined by (8) then $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists for any $p \in F(T)$.

Proof. Let any $p \in F(T)$, using (8) we get

$$\begin{aligned} & \|x_{n+1} - p\| = \|a_n x_n + b_n T^n x_n + c_n R^n x_n + d_n S^n y_n - p\| \\ & = \|a_n (x_n - p) + b_n (T^n x_n - p) + c_n (R^n x_n - p) + d_n (S^n y_n - p)\| \\ & \leq a_n \|x_n - p\| + b_n k_n \|x_n - p\| + c_n k_n \|x_n - p\| + d_n k_n \|y_n - p\| \\ & \leq a_n \|x_n - p\| + b_n k_n \|x_n - p\| + c_n k_n \|x_n - p\| \\ & + d_n k_n (a_n' \|x_n - p\| + b_n' \|S^n x_n - p\| + c_n' \|T^n x_n - p\| + d_n' \|R^n x_n - p\|) \\ & \leq a_n \|x_n - p\| + b_n k_n \|x_n - p\| + c_n \|x_n - p\| + d_n a_n' k_n (\|x_n - p\| + d_n b_n' k_n^2 \|x_n - p\| + d_n c_n' k_n^2 \|x_n - p\| + d_n d_n' k_n^2 \|x_n - p\|) \\ & \leq (1 + b_n (k_n - 1) + c_n (k_n - 1) + d_n (k_n^2 - 1) - d_n a_n' k_n (k_n - 1)) \|x_n - p\| \\ & \leq [1 + \{b_n + c_n + d_n (k_n + 1) - d_n a_n' k_n\} (k_n - 1)] \|x_n - p\| \\ & \leq [1 + \{b_n + c_n + d_n k_n (1 - a_n')\} (k_n - 1)] \|x_n - p\| \\ & \leq [1 + \{1 - a_n + d_n k_n (1 - a_n')\} (k_n - 1)] \|x_n - p\| \end{aligned}$$

Since $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ and $k_n \rightarrow 1$ as $n \rightarrow \infty$, consequently, the condition of Lemma 3 follows from Lemma 1. This completes the proof.

3. MAIN RESULT

In this section, we have proved the approximate common fixed points of three asymptotically nonexpansive mappings for weak and strong convergence results, using iteration process. In the consequence, F denotes the set of common fixed point of the mappings T, R and S .

Theorem-1 Let E be a uniformly convex Banach space and K be a nonempty closed convex subset of E . Let $T, S, R : K \rightarrow K$ be three asymptotically nonexpansive self mappings of K

with $\sum_{n=1}^{\infty} (k_n - 1) < \infty$ satisfying the iteration (8). Let $\{a_n\}, \{b_n\}, \{c_n\}, \{d_n\}, \{a_n'\}, \{b_n'\}, \{c_n'\}, \{d_n'\}$, is in $[0, 1]$ and $a_n + b_n + c_n + d_n = a_n' + b_n' + c_n' + d_n' = 1$. If $F \neq \emptyset$, $k_n \rightarrow 1$ as $n \rightarrow \infty$, then $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = \lim_{n \rightarrow \infty} \|x_n - Rx_n\| = \lim_{n \rightarrow \infty} \|x_n - Sx_n\| = 0$.

Proof. Let $P \in F$. since $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists as proved in Lemma 3. Suppose $\lim_{n \rightarrow \infty} \|x_n - p\| = c$, where $c \geq 0$ is a real number. Now suppose $c > 0$

Thus

$$\begin{aligned} \|y_n - p\| &= \|a_n' x_n + b_n' S^n x_n + c_n' T^n x_n + d_n' R^n x_n - p\| \\ &\leq a_n' \|x_n - p\| + b_n' \|S^n x_n - p\| + c_n' \|T^n x_n - p\| + d_n' \|R^n x_n - p\| \\ &\leq a_n' \|x_n - p\| + b_n' k_n \|x_n - p\| + c_n' k_n \|x_n - p\| + d_n' k_n \|x_n - p\| \\ &\leq (a_n' + b_n' k_n + c_n' k_n + d_n' k_n) \|x_n - p\| \\ &\leq \{1 + a_n'(k_n - 1)\} \|x_n - p\| \\ &\leq \{1 + (1 - c_n')(k_n - 1)\} \|x_n - p\|. \end{aligned}$$

Taking \limsup on both sides of the above inequality, we get

$$\limsup_{n \rightarrow \infty} \|y_n - p\| \leq c \tag{10}$$

Also from

$$\|T^n x_n - p\| \leq k_n \|x_n - p\|$$

For all $n = 1, 2, \dots$, we have

$$\limsup_{n \rightarrow \infty} \|T^n x_n - p\| \leq c \tag{11}$$

Next, we consider

$$\|S^n y_n - p\| \leq k_n \|y_n - p\|$$

Taking \limsup on both sides of the above inequality and using (10), we have

$$\limsup_{n \rightarrow \infty} \|S^n y_n - p\| \leq c$$

Furthermore,

$$c = \lim_{n \rightarrow \infty} \|x_{n+1} - p\|$$

$$\leq \lim_{n \rightarrow \infty} (\leq a_n \|x_n - p\| + b_n k_n \|x_n - p\| + c_n k_n \|x_n - p\| + d_n k_n \|y_n - p\|)$$

$$= (1 - d_n) \|T^n x_n - p\| + d_n \|S^n y_n - p\|$$

By lemma 2 we get

$$\lim_{n \rightarrow \infty} \|T^n x_n - S^n y_n\| = 0 \quad (12)$$

And similarly,

$$\lim_{n \rightarrow \infty} \|R^n x_n - S^n y_n\| = 0 \quad (13)$$

Now,

$$\begin{aligned} \|x_{n+1} - p\| &= \|a_n x_n + b_n T^n x_n + c_n R^n x_n + d_n S^n y_n - p\| \\ &= \|a_n x_n + b_n T^n x_n + c_n x_n + d_n S^n y_n - p\| \\ &= \|a_n x_n + b_n T^n x_n + c_n x_n + d_n S^n y_n - p\| \\ &= \|a_n T^n x_n + b_n T^n x_n + c_n T^n x_n + d_n S^n y_n - p\| \\ &\leq \|(1 - d_n) T^n x_n + d_n S^n y_n - p\| \\ &\leq \|T^n x_n - p\| + d_n \|T^n x_n - S^n y_n\| \end{aligned}$$

which yields that

$$c \leq \liminf_{n \rightarrow \infty} \|T^n x_n - p\|$$

so that (11) gives

$$\lim_{n \rightarrow \infty} \|T^n x_n - p\| = c$$

$$\begin{aligned} \text{In turn, } \|T^n x_n - p\| &\leq \|T^n x_n - S^n y_n\| + \|S^n y_n - p\| \\ &\leq \|T^n x_n - S^n y_n\| + kn \|y_n - p\| \end{aligned}$$

Which implies that

$$c \leq \liminf_{n \rightarrow \infty} \|y_n - p\| \quad (14)$$

By (9) and (12), we have

$$\leq \lim_{n \rightarrow \infty} \|y_n - p\| = c \quad (15)$$

Again, we get

$$c = \lim_{n \rightarrow \infty} \|y_n - p\| = \lim_{n \rightarrow \infty} (1 - d_n) \|x_n - p\| + d_n \|T^n x_n - p\|$$

gives by Lemma 2 that

$$= \lim_{n \rightarrow \infty} \|T^n x_n - x_n\| = 0 \quad (16)$$

Notice that

$$\|y_n - x_n\| = d_n \|T^n x_n - x_n\|$$

Hence by (16)

$$\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0 \quad (17)$$

Now

$$\begin{aligned} \|x_{n+1} - x_n\| &= \|a_n x_n + b_n T^n x_n + c_n R^n x_n + d_n S^n y_n - x_n\| \\ &\leq (1 - a_n) \|T^n x_n - x_n\| + d_n \|T^n x_n - S^n y_n\| + c_n \|T^n x_n - R^n x_n\| \end{aligned}$$

This gives

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0 \quad (18)$$

So that

$$\|x_{n+1} - y_n\| \leq \|x_{n+1} - x_n\| + \|y_n - x_n\| \rightarrow 0 \text{ as } n \rightarrow \infty$$

This gives

$$\lim_{n \rightarrow \infty} \|x_{n+1} - y_n\| = 0. \quad (19)$$

Moreover, from

$$\|x_{n+1} - S^n y_n\| \leq \|x_{n+1} - x_n\| + \|x_n - T^n x_n\| + \|T^n x_n - S^n y_n\|$$

Which gives that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - S^n y_n\| = 0 \quad (20)$$

Using (12),(16)and(17) we obtained

$$\begin{aligned} \|x_n - S^n x_n\| &\leq \|x_n - T^n x_n\| + \|T^n x_n - S^n y_n\| + \|S^n y_n - S x_n\| \\ &\leq \|x_n - T^n x_n\| + \|T^n x_n - S^n y_n\| + k \|y_n - x_n\| \end{aligned}$$

Gives that

$$\lim_{n \rightarrow \infty} \|x_n - S^n x_n\| = 0$$

And

$$\begin{aligned} \|x_{n+1} - S x_{n+1}\| &\leq \|x_{n+1} - S^{n+1} x_{n+1}\| + \|S^{n+1} x_{n+1} - S x_{n+1}\| \\ &\leq \|x_{n+1} - S^{n+1} x_{n+1}\| + k \|S^{n+1} x_{n+1} - S x_{n+1}\| \\ &\leq \|x_{n+1} - S^{n+1} x_{n+1}\| + k (\|S^n x_{n+1} - S^n y_n\| + k \|S^n y_n - x_{n+1}\|) \\ &\leq \|x_{n+1} - S^{n+1} x_{n+1}\| + k^2 \|x_{n+1} - y_n\| + k \|S^n y_n - x_{n+1}\| \end{aligned}$$

This implies that

$$\lim_{n \rightarrow \infty} \|x_n - Sx_n\| = 0$$

Now

$$\begin{aligned} \|x_{n+1} - Tx_{n+1}\| &\leq \|x_{n+1} - T^{n+1}x_{n+1}\| + \|T^{n+1}x_{n+1} - T^{n+1}x_n\| + \|T^{n+1}x_n - Tx_{n+1}\| \\ &\leq \|x_{n+1} - T^{n+1}x_{n+1}\| + k \|x_{n+1} - x_n\| + k \|T^n x_n - x_{n+1}\| \\ &= \|x_{n+1} - T^{n+1}x_{n+1}\| + k \|x_{n+1} - x_n\| + k a_n \|T^n x_n - S^n y_n\| \end{aligned}$$

Which yields

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$$

Similarly,

This completes the proof.

$$\lim_{n \rightarrow \infty} \|x_n - Rx_n\| = 0$$

Theorem-2 Let E be a uniformly convex Banach space satisfying Opial condition and K, T, S and $\{x_n\}$ be taken as theorem 1. If $F(S) \cap F(T) \cap F(R) \neq \emptyset$, $I - T$, $I - R$ and $I - S$ are demiclosed, at zero, then $\{x_n\}$ converges weakly to a common fixed point of S, R and T.

Proof. Let $p \in F(S) \cap F(T) \cap F(R)$. Then as proved in Lemma 3 $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists. Since E is uniformly convex. Thus there exists subsequences $\{x_{n_k}\}$ of $\{x_n\}$ such that $\{x_{n_k}\}$ converges weakly to $z_1 \in K$. From theorem 1, we have

$$\lim_{n \rightarrow \infty} \|x_{n_k} - Tx_{n_k}\| = 0, \lim_{n \rightarrow \infty} \|x_{n_k} - Sx_{n_k}\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|x_{n_k} - Rx_{n_k}\| = 0$$

Since $I - T$, $I - R$ and $I - S$ are demiclosed, at zero, therefore $Sz_1 = z_1$. Similarly $Tz_1 = z_1$ and $Rz_1 = z_1$. Again in the same way, we can prove that $z_2 \in F(S) \cap F(T) \cap F(R)$. Next, we prove the uniqueness. From Lemma 3 the limits $\lim_{n \rightarrow \infty} \|x_n - z_2\|$ exists. For this suppose that $z_1 \neq z_2$, then by the Opial's condition

$$\lim_{n \rightarrow \infty} \|x_n - z_1\| = \lim_{n_i \rightarrow \infty} \|x_{n_i} - z_1\| < \lim_{n_i \rightarrow \infty} \|x_{n_i} - z_2\|$$

$$= \lim_{n \rightarrow \infty} \|x_n - z_2\| = \lim_{n_j \rightarrow \infty} \|x_{n_j} - z_2\|$$

$$< \lim_{n_j \rightarrow \infty} \|x_{n_j} - z_1\| < \lim_{n \rightarrow \infty} \|x_n - z_1\|.$$

This is a contradiction so $z_1 = z_2$, Hence $\{x_n\}$ converges weakly to a common fixed point of T, R and S.

Theorem-3 Let E be a real uniformly convex Banach space and K, S, R, T, F, $\{x_n\}$ be as in theorem 1. Then $\{x_n\}$ converges strongly to a common fixed point of T, R and S if and only if

$$\liminf_{n \rightarrow \infty} d(x_n, F) = 0$$

Proof. Necessity is evident, let $\lim_{n \rightarrow \infty} d(x_n, F) = 0$. From Lemma 3, $\lim_{n \rightarrow \infty} \|x_n - p\|$ exists for all $p \in F$, so that $\lim_{n \rightarrow \infty} d(x_n, F)$ exists. Since by hypothesis, $\liminf_{n \rightarrow \infty} d(x_n, F) = 0$, so that, we get

$$\lim_{n \rightarrow \infty} d(x_n, F) = 0.$$

But $\{x_n\}$ is Cauchy sequence and therefore converges to p . We know that $\lim_{n \rightarrow \infty} d(x_n, F) = 0$, we obtained $d(p, F) = 0$, therefore $p \in F$. Using theorem 3, we obtain a strong convergence theorem of the iteration scheme (8) under the Condition (A').

Theorem-4 Let E be a uniformly convex Banach space and $K, S, R, T, F, \{x_n\}$ be as in Theorem 3. Let S, T, R satisfy the Condition (A') and $F \neq \emptyset$. Then $\{x_n\}$ converges strongly to a point of F .

Proof . We proved in Theorem 5, i.e.

$$\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$$

or

$$\lim_{n \rightarrow \infty} \|x_n - Rx_n\| = 0$$

or

$$\lim_{n \rightarrow \infty} \|x_n - Sx_n\| = 0$$

Then from the definition of Condition (A'), we obtain

$$\lim_{n \rightarrow \infty} f(d(x_n, F)) \leq \lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$$

or

$$\lim_{n \rightarrow \infty} f(d(x_n, F)) \leq \lim_{n \rightarrow \infty} \|x_n - Sx_n\| = 0$$

or

$$\lim_{n \rightarrow \infty} f(d(x_n, F)) \leq \lim_{n \rightarrow \infty} \|x_n - Rx_n\| = 0$$

In above cases, we get

$$\lim_{n \rightarrow \infty} f(d(x_n, F)) = 0.$$

But $f : [0, \infty) \rightarrow [0, \infty)$ is a nondecreasing function satisfying $f(0) = 0$, $f(r) > 0$ for all $r \in (0, \infty)$, so that we get $\lim_{n \rightarrow \infty} f(d(x_n, F)) = 0$. All the condition of theorem 3 are satisfied, therefore by its conclusion $\{x_n\}$ converges strongly to a fixed point of F .

Corollary 5. Let K be a nonempty closed convex subset of a uniformly convex Banach space E . Suppose T be a asymptotically nonexpansive mapping of K . Let $\{x_n\}$ be defined by the iteration (8), where $\{d_n\}$ and $\{b'_n\}$ in $[0, 1]$ for all $n \in \mathbb{N}$, then $\{x_n\}$ converges strongly to a fixed point of T .

Proof. Suppose $T = I$ in the above theorem.

Corollary 6. Suppose E be a Banach space satisfying Opial condition and let K and T be taken as in (5). Let $F(T) \neq \emptyset$. Now if the mapping $I - T$, is demiclosed, at zero, then $\{x_n\}$ defined by (3) converges weakly to a fixed point of T .

Corollary 7 Let E be a uniformly convex Banach space which has a Frechet differentiable norm and let K and T be taken as theorem 3. Let $F(T) \neq \emptyset$ taken $\{x_n\}$ defined by (8) converges weakly to a fixed point of T .

Corollary 8 Suppose E be a Banach space satisfying Opial condition and let K and T be taken as in (5). Let $F(T) \neq \emptyset$. Now if the mapping $I - T$, is demiclosed, at zero, then $\{x_n\}$ defined by (8) converges weakly to a fixed point of T .

4. REFERENCES

- [1] B.L. Xu and M.A. Noor, Fixed point iterations for asymptotically non-expansive mappings in Banach spaces, *J. Math. Anal. Appl.* 267 (2002), 444-453.
- [2] G. Das and J.P. Debta, Fixed points of quasi-nonexpansive mappings, *Indian J. Pure. Appl. Math.* (1986), 1263-1269.
- [3] H. Fukhar-ud-din and S.H. Khan, Convergence of iterate with errors of asymptotically quasi-nonexpansive mappings and application, *J. Math. Anal. Appl.* 328 (2007), 821-829.
- [4] H.K. Xu, Existence and convergence for fixed points of asymptotically nonexpansive type, *Nonlinear Anal.* 16 (1991), 1139-1146.
- [5] H.K. Xu and R.G. Ori, An implicit iteration process for nonexpansive mappings, *Numeric Functional Analysis and optimization*, 22(5-6)(2001),707-773.
- [6] J. Schu, Weak and strong convergence to fixed points of asymptotically nonexpansive mappings, *Bull. Aust. Math. Soc.* 43 (1991), 153-159.
- [7] J. Schu, Iterative constructions of fixed points of asymptotically nonexpansive mappings, *J. Math. Anal. Appl.* 158 (1991), 407-413.
- [8] Jing, Zhao, He Songnian and Su, Yongfu, Weak and strong convergence theorems for nonexpansive mapping in Banach space, *Indawi Publishing corporation fixed point theory and application volume 2008*, article ID751383, 7 pages doi:10.1155/2008/751383.
- [9] K. Goebel and W.A. Kirk, A fixed point theorem for asymptotically nonexpansive mappings, *Proc. Amer. Math. Soc.* 35 (1972), 171-174.
- [10] K.K. Tan and H.K. Xu, Approximating fixed point of nonexpansive mappings by Ishikawa iteration process, *J. Math. Anal. Appl.* 178 (1993), 301-308.
- [11] K. Nammanee, M.A. Noor and S. Suantai, Convergence criteria of modifies Noor iteration with errors for asymptotically nonexpansive mappings, *J.Math. Anal. Appl.* 4(2006), 320-334.
- [12] L.C. Deng, P. Cubiotti and J.C. Yao, Approximation of common fixed points of families of nonexpansive mappings, *Taiwanese J. Math.* 12(2008), 487-500.
- [13] L.C. Deng, P. Cubiotti, and J.C. Yao, An implicit iteration scheme for monotone variation inequalities and fixed point problems, *Nonlinear Anal.* 69 (2008), 2445-2457.
- [14] L.C. Deng, S. Schaible and J.C. Yao, Implicit iteration scheme with per-tarbed mapping for equilibrium problems and fixed point problems offinetely many nonexpansive mapping, *J. Optim.Theory Appl.* 139 (2008), 403-418.
- [15] M.R. Yadav and B.P. Tripathi, Iteration scheme for common fixed points of two asymptotically nonexpansive mappings, *International Journal of Pure and Applied Mathematics* 80(4)(2012),501-514.
- [16] Q.H. Liu, Iteration sequences for asymptotically quasi-nonexpansive mapping with an error member in a uniformly convex Banach space, *J. Math. Anal. Appl.* 266 (2002), 468-471.
- [17] R.P. Agrawal, Donal O'Regan and D.R. Sahu, Iterative construction offixed points; of nearly asymptotically nonexpansive mappings, *J. Nonlinear Convex, Anal.* 8(1) (2007) 61-79
- [18] S.H. Khan, Y.J. Cho and M. Abbas, Convergence to common fixed points by a modified iteration process, *J. Appl. Math. And Comput.*doi:10.1007/s12190-010-0381.
- [19] S.H. Khan and W. Takahashi, Approximating common fixed points of two asymptotically nonexpansive mappings, *Sci. Math. Jpn.* 53(1) (2001), 143-148.
- [20] W.R. Mann, Mean value methods in iteration, *Proc. Amer. Math. Soc.* 4(1953), 506-510.
- [21] W. Takahashi, and G.E. Kim, Approximating fixed points of nonexpansive mappings in Banach spaces, *Math. Japon.* 48(1) (1998), 1-9.
- [22] W. Nisrakoo and S. Saejung, A new three-step fixed point iteration scheme for asymptotically nonexpansive mappings, *Comput. Math. Appl.*18 (2006), 1026-1034
- [23] W. Takahashi and T. Tamura, Limit theorems of operators by convex combinations of nonexpansive retractions in Banach spaces, *J. Approx.Theory*, 91(3) (1997), 386-397.
- [24] W. Takahashi, Iterative methods for approximation of fixed points and their applications, *J. Oper. Res. Soc. Jpn.* 43(1) (2000), 87-108.
- [25] Y.J. Cho, H.Y. Zhou and G. Guo, Weak and strong convergence theorems for three-step iterations with errors for asymptotically nonexpansive mapping, *Comput. Math. Appl.*, 47 (2004), 707-717.
- [26] Z. Opial, Weak and convergence of successive approximations for nonexpansive mapping, *Bull. Amer. Math. Soc.* 73 (1967), 591-597.